

Freshwater sponge References

- Adams, E., G. Goss and S. Leys (2010). "Freshwater Sponges Have Functional, Sealing Epithelia with High Transepithelial Resistance and Negative Transepithelial Potential." Public Library of Science One **5**.
- Adams, E. D. M. (2010). Physiology and morphology of epithelia in the freshwater demosponge, *Spongilla lacustris*. MSc, University of Alberta.
- Annandale, N. (1971). "Notes on freshwater sponges. VI. The midday siesta of *Spongilla* in the tropics." Records of the Indian Museum : a journal of Indian zoology **1**: 387-392.
- Bannister, R. J., C. N. Battershill and R. de Nys (2010). "Demographic variability and long-term change in a coral reef sponge along a cross-shelf gradient of the Great Barrier Reef." Marine and Freshwater Research **61**(4): 389-396.
- Bergquist, P. R. and M. E. Sinclair (1968). "The morphology and behaviour of larvae of some intertidal sponges." New Zealand Journal of Marine and Freshwater Research **2**: 426-437.
- Bond, C. (1992). "Continuous cell movements rearrange anatomical structures in intact sponges." Journal of Experimental Zoology **263**: 284-302.
- Bond, C. and A. K. Harris (1988). "Locomotion of sponges and its physical mechanism." Journal of Experimental Zoology **246**: 271-284.
- Coutinho, C., J. Seack, G. Van de Vyver, R. Borojevic and W. E. G. Müller (1998). "Origin of the Metazoan Bodyplan: Characterization and Functional Testing of the Promoter of the Homeobox Gene *EmH-3* from the Freshwater Sponge *Ephydatia muelleri* in Mouse 3T3 Cells." Biological Chemistry **379**: 1243-1252.
- Coutinho, C. C., J. Seack, G. Van de Vyver, R. Borojevic and W. E. G. Mueller (1994). "Origin of the metazoan bodyplan: characterization and functional testing of the promoter of the homeobox gene *EmH-3* from the freshwater sponge *Ephydatia muelleri* in Mouse 3T3 cells." Biology and Chemistry **379**: 1243-1251.
- Covich, A. P., M. C. Austen, F. BÄRlocher, E. Chauvet, B. J. Cardinale, C. L. Biles, P. Inchausti, O. Dangles, M. Solan, M. O. Gessner, B. Statzner and B. Moss (2004). "The Role of Biodiversity in the Functioning of Freshwater and Marine Benthic Ecosystems." BioScience **54**(8): 767-775.
- Drum, R. W. (1968). "Electron microscopy of siliceous spicules from the freshwater sponge *Heteromyenia*." Journal of Ultrastructure Research **22**: 12-21.
- Elliott, G. and S. Leys (2007). "Coordinated behaviour in a freshwater sponge: Orchestrated contractions effectively expel water from the aquiferous system." Journal of Experimental Biology **210**: 3736-3748.
- Elliott, G. R. D. (2009). The cytology and physiology of coordinated behaviour in a freshwater sponge, *Ephydatia muelleri*. PhD, University of Alberta.
- Elliott, G. R. D. and S. P. Leys (2003). "Sponge coughing: Stimulated contractions in a juvenile freshwater sponge, *Ephydatia muelleri*." Integrative and Comparative Biology **43**: 817-817.
- Elliott, G. R. D. and S. P. Leys (2010). "Evidence for glutamate, GABA and NO in coordinating behaviour in the sponge, *Ephydatia muelleri* (Demospongiae, Spongillidae)." Journal of Experimental Biology **213**(13): 2310-2321.
- Frost, T. M. (1980). "Clearance rate determinations for the freshwater sponge *Spongilla lacustris*: effects of temperature, particle type and concentration, and sponge size." Archives fur Hydrobiologie **90**(3): 330-356.

- Funayama, N., M. Nakatsukasa, S. Kuraku, K. Takechi, M. Dohi, N. Iwabe, T. Miyata and K. Agata (2005). "Isolation of *Ef silicatein* and *Ef lectin* as molecular markers of sclerocytes and cells involved in innate immunity in the freshwater sponge *Ephydatia fluviatilis*." Zoological Science **22**(10): 1113-1122.
- Harrison, F., N. Kaye and G. Kaye (1985). The dermal membrane of *Eunapius fragilis*. New Perspectives in Sponge Biology, Woods Hole, MA, Smithsonian Institution Press.
- Harrison, F. W. (1972). "The nature and role of the basal pinacoderm of *Corvomeyenia carolinensis* Harrison (Porifera: Spongillidae)." Hydrobiologia **39**(4): 495-508.
- Holmes, R. M., A. Aminot, R. K  rouel, B. A. Hooker and B. J. Peterson (1999). "A simple and precise method for measuring ammonium in marine and freshwater ecosystems." Canadian Journal of Fisheries and Aquatic Sciences **56**(10): 1801-1808.
- Huysecom, J., E. Richelle-Maurer, G. Van de Vyver and B. Vray (1988). "Effect of bacterial concentration on retention and growth rate of the freshwater sponge *Ephydatia fluviatilis*." Physiological Zoology **61**(6): 535-542.
- Insiecke, G. (1994). Ingestion and digestion of *Chlamydomonas reinhardtii* (Volvocales) by the freshwater sponge *Spongilla lacustris* (Spongillidae). Sponges in Time and Space. R. W. M. van Soest, T. M. G. van Kempen and J.-C. Braekman. Rotterdam, A.A. Balkema: 371-376.
- Insiecke, G., R. Borojevic, M. Custodio and W. E. G. M  ller (1994). "Retinoic acid acts as a morphogen in freshwater sponges." Invertebrate Reproduction and Development **26**(2): 89-98.
- Insiecke, G., U. Pascheberg and W. E. G. M  ller (1993). "Preparation and karyotype analysis of mitotic chromosomes of the freshwater sponge *Spongilla lacustris*." Chromosoma **102**: 724-727.
- Insiecke, G., R. Steffen, M. Custodio, R. Borojevic and W. E. G. M  ller (1995). "Formation of spicules by sclerocytes from the freshwater sponge *Ephydatia muelleri* in short-term cultures in vitro." In vitro Cellular and Developmental Biology-Animal **31**: 528-535.
- Karpov, S. A. and B. S. C. Leadbeater (1997). "Cell and Nuclear Division in a Freshwater Choanoflagellate, *Monosiga ovata* Kent." European Journal of Protistology **33**: 323-334.
- Koonce, M. P., U. Euteneuer, K. L. McDonald, D. Menzel and M. Schliwa (1986). "Cytoskeletal architecture and motility in a giant freshwater amoeba, *Reticulomyxa*." Cell Motility and the Cytoskeleton **6**: 521-533.
- Koonce, M. P. and M. Schliwa (1986). "Reactivation of organelle movements along the cytoskeletal framework of a giant freshwater amoeba." Journal of Cell Biology **103**: 605-612.
- Langenbruch, P. F. and N. Weissenfels (1987). "Canal systems and choanocyte chambers in freshwater sponges (Porifera, Spongillidae)." Zoomorphology **107**: 11-16.
- Mackie, G. O. (1999). "Coelenterate Organs." Marine and Freshwater Behavioral Physiology **32**: 113-127.
- Meixner, M. J., C. L  ter, C. Eckert, V. Itskovich, D. Janussen, T. von Rintelen, A. V. Bohne, J. M. Meixner and W. R. Hess (2007). "Phylogenetic analysis of freshwater sponges provide evidence for endemism and radiation in ancient lakes." Molecular Phylogenetics and Evolution **45**(3): 875-886.
- Mohri, K., M. Nakatsukasa, Y. Masuda, K. Agata and N. Funayama (2008). "Toward understanding the morphogenesis of siliceous spicules in freshwater sponge: differential

- mRNA expression of spicule-type-specific silicatein genes in *Ephydatia fluviatilis*." Developmental dynamics **237**: 3024-3039.
- Mukai, H. (1989). "Growth and reproduction of four species of freshwater sponge cultured in their natural surroundings." Science Reports of the Faculty of Education, Gunma University **38**: 25-47.
- Mukai, H. (1990). "Further studies on growth and sex differentiation in four species of freshwater sponges." Science Reports of the Faculty of Education, Gunma University **39**: 41-56.
- Nikko, E., G. Van de Vyver and E. Richelle-Maurer (2001). "Retinoic acid down-regulates the expression of EmH-3 homeobox-containing gene in the freshwater sponge *Ephydatia muelleri*." Mechanisms of Ageing and Development **122**: 779-794.
- Rader, R. B. and R. N. Winget (1985). "Seasonal growth rate and population dynamics of a freshwater sponge." Hydrobiologia **123**: 171-176.
- Rasmont, R., J. Bouillon, P. Castiaux and G. Vandermeerssche (1958). "Ultra-structure of the choanocyte collar cells in freshwater sponges." Nature **181**: 58-59.
- Richelle-Maurer, E. and G. Van de Vyver (1999). "Temporal and spatial expression of *EmH-3*, a homeobox-containing gene isolated from the freshwater sponge *Ephydatia muelleri*." Mechanisms of Ageing and Development **109**: 203-219.
- Richelle-Maurer, E., G. Van de Vyver, S. Vissers and C. C. Coutinho (1998). Homeobox-containing genes in freshwater sponges: characterization, expression, and phylogeny. Molecular Evolution: Evidence for Monophyly of Metazoa. W. E. G. Müller. Berlin, Heidelberg, Springer Berlin Heidelberg. **19**: 157-175.
- Rivera, A. S., J. U. Hammel, K. M. Haen, E. S. Danka, B. Cieniewicz, I. P. Winters, D. Posfai, G. Wörheide, D. V. Lavrov, S. W. Knight, M. S. Hill, A. L. Hill and M. Nickel (2011). "RNA interference in marine and freshwater sponges: actin knockdown in *Tethya wilhelma* and *Ephydatia muelleri* by ingested dsRNA expressing bacteria." BMC Biotechnology **11**: 67.
- Schierenberg, E. (2005). "Unusual cleavage and gastrulation in a freshwater nematode: developmental and phylogenetic implications." Developmental Genes and Evolution **215**: 103-108.
- Schroder, H. C., S. M. Efremova, V. B. Itskovich, S. Belikov, Y. Masuda, A. Krasko, I. M. Muller and W. E. G. Muller (2002). "Molecular Phylogeny of the freshwater sponges in Lake Baikal." J. Zool. Syst. Evol. Research **40**: 1-7.
- Seimiya, M., Y. Watanabe and Y. Kurosawa (1997). "Identification of POU-class homeobox genes in a freshwater sponge and the specific expression of these genes during differentiation." European Journal of Biochemistry **243**: 27-31.
- Simpson, T. L., R. Garrone and M. Mazzorana (1983). "Interaction of germanium with biosilicification in the freshwater sponge *Ephydatia mülleri*: evidence of localized membrane domains in the silicalemma." Journal of Ultrastructure Research **85**: 159-174.
- Simpson, T. L. and C. A. Vaccaro (1974). "An Ultrastructural Study of Silica Deposition in the Freshwater Sponge *Spongilla lacustris*." Journal of ultrastructure research **47**: 296-309.
- Strekal, T. A. and W. McDiffett (1974). "Factors affecting germination, growth, and distribution of the freshwater sponge, *Spongilla fragilis* Leidy (Porifera)." Biological Bulletin **146**: 267-278.

- Wachtmann, D., W. Stockem and N. Weissenfels (1990). "Cytoskeletal organization and cell organelle transport in basal epithelial cells of the freshwater sponge *Spongilla lacustris*." Cell and Tissue Research **261**: 145-154.
- Weissenfels, N. (1992). "The filtration apparatus for food collection in freshwater sponges (Porifera, Spongillidae)." Zoomorphology **112**: 51-55.
- Wilkinson, C. R. (1979). "Skeletal morphology of coral reef sponges: A scanning electron microscope study." Australian Journal of Marine and Freshwater Research **30**: 793-801.
- Willenz, P. (1980). Kinetic and morphological aspects of the particle ingestion by the freshwater sponge *Ephydatia fluviatilis*. Nutrition in the Lower Metazoa. D. C. Smith and Y. Tiffon. Oxford, Pergamon Press: 163-178.
- Willenz, P. and G. Van de Vyver (1986). "Ultrastructural evidence of extruding exocytosis of residual bodies in the freshwater sponge *Ephydatia fluviatilis*." Journal of Morphology **190**: 307-318.
- Willenz, P., B. Vray, M. P. Maillard and G. Van de Vyver (1986). "A quantitative study of the retention of radioactively labeled *E. coli* by the freshwater sponge *Ephydatia fluviatilis*." Physiological and Biochemical Zoology **59**(5): 495-504.
- Windsor, P. J. (2014). Evolution of the sponge body plan: Wnt and the development of polarity in freshwater sponges. PhD Doctoral Dissertation, University of Alberta.